

## RESEARCH ARTICLE OPEN ACCESS

# Sun Bear Ecology and Conservation: Multiscale Habitat Associations in Southeast Asia

Alexander Hendry<sup>1</sup> | Harsh K. Pahuja<sup>2</sup> | Zachary Amir<sup>3,4,5</sup> | Ilyas Nursamsi<sup>1</sup> | Jonathan H. Moore<sup>6</sup> | Matthew Scott Luskin<sup>3,5</sup> 

<sup>1</sup>School of the Environment, University of Queensland, Brisbane, Queensland, Australia | <sup>2</sup>School of Agriculture and Food Sciences, University of Queensland, Brisbane, Queensland, Australia | <sup>3</sup>School of the Environment & Centre for Biodiversity and Conservation Science, University of Queensland, Brisbane, Queensland, Australia | <sup>4</sup>Terrestrial Ecosystem Research Network, University of Queensland, Brisbane, Queensland, Australia | <sup>5</sup>The Wildlife Observatory of Australia (WildObs), QCIF, Brisbane, Queensland, Australia | <sup>6</sup>Asia Pacific, Fauna and Flora International, Phnom Penh, Cambodia

**Correspondence:** Matthew Scott Luskin ([m.luskin@uq.edu.au](mailto:m.luskin@uq.edu.au))

**Received:** 10 February 2025 | **Accepted:** 9 January 2026

**Editor-in-Chief & Handling Editor:** Ahimsa Campos-Arceiz

**Funding:** Smithsonian Institution's Forest Global Earth Observatory (ForestGEO); National Geographic Society, Grant/Award Number: 9384-13; Australian Research Council, Grant/Award Number: DE21010144

**Keywords:** camera trapping | diel activity | *Helarctos malayanus* | oil palm plantation | rainforest logging

## ABSTRACT

Degraded tropical forests, defined here as logged, fragmented, and edge forests, constitute most of the remaining natural areas in Southeast Asia, potentially contributing to megafauna declines and extirpations. The sun bear (*Helarctos malayanus*) is a megafauna species native to Southeast Asian forests and is generally considered a forest specialist, and thus may be at risk from widespread forest degradation. To clarify the sun bear's responses to forest degradation, we assessed landscape- and local-level occurrence and diel activity across 38 landscapes throughout its range. First, we assess among-landscape variation in sun bear detections by comparing detection records from 185 camera studies. Second, we assess within-landscape variation in abundance using 19 new camera surveys in eight landscapes and Royle–Nichols hierarchical modelling to account for imperfect detectability. Finally, we assessed diel activity in intact versus degraded forests. Across the region (among landscapes), sun bear detections were higher in larger forests, but unexpectedly also in forests that neighbour humans. Within landscapes (i.e., local-scale), there were no significant responses to forest degradation but sun bears avoided oil palm plantations and had positive associations with elevation and humans. The diel activity analysis showed that sun bears are predominantly crepuscular and do not shift their activity patterns in degraded forests or near humans. These findings challenge prior work suggesting sun bears are specialists and suggest they are more tolerant of humans and forest degradation than previously realised. The sun bear's adaptability may be traced to its being a habitat generalist and omnivore, rather than to temporal partitioning to avoid humans. Our multi-scale synthesis suggests that, where hunting is low, degraded forests contribute to sun bear conservation.

## 1 | Introduction

Globally, megafauna, usually defined as terrestrial mammals with an average adult mass greater than 40 kg, are experiencing significant population declines (Harrison et al. 2016; Ripple et al. 2016, 2019). Southeast Asia, in particular, is suffering severe

losses of megafauna in recent decades (Ripple et al. 2016; Mahmood et al. 2021; Amir et al. 2022), largely due to the widespread poaching (especially snaring), as well as deforestation and forest degradation (defined as logged, fragmented, and edge forests) (Hughes 2017; Strang and Rusli 2021). One Southeast Asian

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2026 The Author(s). *Integrative Conservation* published by John Wiley & Sons Australia, Ltd on behalf of Xishuangbanna Tropical Botanical Garden (XTBG).

## Summary

Much of the natural environment in Southeast Asia has been degraded by humans, and along with hunting, these factors often contribute to declines of megafauna, such as sun bears (*Helarctos malayanus*). To determine the sun bear's response to forest degradation, we assessed sun bear detections at two spatial scales, regional (landscape-level) and locally (within a single landscape). We collated new and previously published camera trap studies and assessed how sun bear detection rates varied with environmental and anthropogenic factors. We also assessed the times at which sun bears were detected to determine whether they change their behaviour in disturbed habitats. At the regional/landscape scale, sun bears were detected most frequently in large forests but were also common in forests that neighboured humans. At the local scale, sun bears were associated with higher elevations and avoided oil palm plantations. We found that sun bears were mainly crepuscular and did not change their behaviour to avoid humans. These results suggest sun bears may be able to survive in areas with low-intensity forest degradation and although the reasons are not clear, it could be due to their omnivorous diet, and not by avoiding humans. Our findings suggest that some moderately degraded forests have high conservation value for sun bears.

### • Practitioner Points

- Across their range, sun bears are associated with large forests but are also present in forests near humans and infrastructure like roads.
- Sun bears are predominantly crepuscular and do not shift their activity pattern to avoid humans.
- Moderately degraded habitat (logged forests and forest edges) has significant conservation value for sun bears and should therefore be protected and managed to reduce hunting.

megafauna species that is threatened by these processes is the world's smallest bear species, the sun bear (*Helarctos malayanus*) (Meijaard 1999; Scotson et al. 2017a).

The sun bear's range extends from Yunnan Province in China southward through continental Southeast Asia and into Sumatra and Borneo (Linkie et al. 2007; Fredriksson 2012; Nazeri et al. 2014; Li et al. 2017). The species has an International Union for the Conservation of Nature Red List (IUCN-RL) classification of "Vulnerable" (Scotson et al. 2017a), as its population is declining due to acute and synergetic threats (Wilcove et al. 2013; Gallego-Zamorano et al. 2020). First, despite being listed in the Convention on the International Trade in Endangered Species (CITES) Supporting Information: Appendix I, sun bears are targeted by hunters and poachers for bushmeat, body parts such as claws for souvenirs, and especially gall bladders for bile used in traditional Chinese medicine (Shepherd and Shepherd 2010; Krishnasamy and Shepherd 2014; Gomez et al. 2020). Sun bear cubs are also targeted for the illegal pet trade (Scheffers et al. 2019). Second, Southeast Asian forests have undergone profound environmental changes in recent decades. There has been extensive land clearing for timber, as well as for crops such

as rubber and oil palm. Additionally, over 70% of Asia's remaining forest is degraded by being within 1 km of an edge and is often selectively logged (Haddad et al. 2015; Grantham et al. 2020). As land clearing often requires the simultaneous construction of roads that provide access to poachers, these two key threats to sun bears (forest degradation and hunting) often interact synergistically (Laurance et al. 2009; Clements et al. 2014). Therefore, the sun bear's continued persistence largely depends on the degree of forest degradation it can tolerate and low hunting.

Sun bears are often considered forest specialists that predominantly occupy intact, undisturbed, lowland, dipterocarp tropical forests (Augeri 2005; Wong et al. 2013; Nazeri et al. 2014). This paradigm has been challenged by studies demonstrating that sun bears can persist in regenerating forests previously degraded by logging, provided poaching is low (Linkie et al. 2007; Brodie et al. 2015; Guharajan et al. 2021; Hwang et al. 2021; Guharajan et al. 2022). Our work follows from that of Guharajan et al. (2021, 2022), and Hwang et al. (2021)—who identified that sun bears can persist in recently or currently logged forests. We assess this by analysing sun bear camera detections relative to forest and anthropogenic covariates on a larger scale. Guharajan et al.'s (2021, 2022), and Hwang et al.'s (2021) work were confined to relatively small geographic areas and may be subject to site-specificity, while our datasets cover a larger portion of the species' range and utilises data sourced from multiple countries. Finally, we conducted analyses at two spatial scales (within and among landscapes), which is informative to managers who operate across different scales, from park rangers working in a single landscape up to regional, national, or international level planning.

Anthropogenic activity may not only reduce the habitat available to sun bears but may also affect their behaviour. Many diurnal and crepuscular animals are known to adjust their behaviour in areas with humans by becoming more nocturnal in order to avoid times of the day when humans are most active (Gaynor et al. 2018; Negret et al. 2023; Lee et al. 2024). Disturbed forests are likely to contain humans, such as people hunting or collecting non-timber forest products like firewood (Fotang et al. 2021), so measures of degradation can be used as a proxy for human presence. However, it is important to acknowledge that forest degradation is not a perfect proxy for human activity, and poachers may focus on intact forests (O'Kelly et al. 2018). Research on sun bear diel activity is limited, with studies from intact forests suggesting that they are predominately diurnal (Te Wong et al. 2004; Augeri 2005; Gouda et al. 2020) and possibly more nocturnal closer to forest edges (Griffiths and Schaik 1993; Tee et al. 2021). This shift to nocturnality could potentially allow sun bears to avoid humans during daylight hours when they are more likely to be present in the landscape (Lee et al. 2024). However, we did not identify any research that suggests that sun bears adapt their diel activity near humans or in degraded forests.

With the exception of Scotson et al. (2017b) and Guharajan et al. (2022), the latter only utilising data from Sabah, Malaysian Borneo, there is a dearth of range-wide ecological research, syntheses, or comparative analyses for sun bears, with the majority of studies focused on small geographic areas (Linkie et al. 2007; Wong and Linkie 2013; Guharajan 2016; Lee et al. 2019). We expand on these earlier studies by including a larger dataset and analysing diel activity alongside habitat

associations. In this study, we examine relationships between sun bear occurrence records collected from several sources across multiple landscapes in the species' range alongside anthropogenic and habitat variables at landscape and local scales. We expected variation in the strength of relationships with the same covariate across the two spatial scales, but that the primary drivers and directions of effects would be consistent. We predicted that sun bears would be positively associated with habitat covariates, including forest size, forest quality, and the Forest Landscape Integrity Index. Likewise, we predicted they would be negatively associated with anthropogenic covariates, including human population, night lights, the Human Footprint Index, and oil palm plantations. This last prediction was less certain as sun bears, like many other animals, are known to enter oil palm plantations where they consume both fruit and insects (Normua et al. 2004; Abidin et al. 2018; Guharajan et al. 2018). However, sun bears may be vulnerable to snaring in plantations, so oil palm and nearby forests may be population sinks (Luskin and Potts 2011; Luskin et al. 2014; Luskin et al. 2018), and they still require neighbouring forests to retreat (Cheah 2013; Guharajan et al. 2018). We therefore predicted that, overall, the extent of oil palm plantations would negatively affect sun bear detections. Finally, we hypothesised that sun bears would shift their diel activity towards nocturnality near humans and in degraded forests, consistent with other species' responses (Lee et al. 2024).

## 2 | Methods

This paper is modelled after several previously published papers (Dehaut et al. 2022; Dunn et al. 2022; Carr et al. 2023; Deceur et al. 2023; Hendry et al. 2023; Honda et al. 2023; Luskin et al. 2023a; Nursamsi et al. 2023; Mendes et al. 2024b), hence the similarities in many parts of the methods.

### 2.1 | Study Region, Data Sources, and Approach

We defined the region of focus for this study as the tropical forest range of the sun bear in Southeast Asia, which specifically includes Myanmar, Thailand, Cambodia, Laos, Peninsular Malaysia, Sumatra, and Borneo [the species has been extirpated from Singapore for over 75 years; (Amir et al. 2022; Brodie et al. 2023; Sartor et al. 2024)]. While sun bears also occur in South Asia (i.e., Northeast India) and infrequently occur in Yunnan province in China, we limited our analysis to the tropical rainforests of Southeast Asia. To make inferences at different scales, we compiled presence and absence data for the sun bear from two sources: (1) survey-level total detections reported in published camera trapping studies and (2) camera-level detection histories from new camera trapping sessions conducted across eight landscapes in Southeast Asia. Presence data consisted of georeferenced occurrence records, defined as the coordinates of a location where a sun bear was observed. We defined a camera study as a continuous sampling effort using at least 10 cameras within a landscape (10–1000 km<sup>2</sup>). We refer to the sampling area as a “landscape”—usually a national park, a production forest, or a collection of nearby forest patches within a 20-km radius area (1256 km<sup>2</sup>). When comparing landscape-level surveys, camera sampling likely reflects population-level

metrics, since individual sun bears have relatively small home ranges compared to other bear species (approx. 15 km<sup>2</sup> from one study—Te Wong et al. 2004). On the contrary, detection histories from landscapes partitioned into many 10 km<sup>2</sup> grid cells likely reflect individual-level habitat associations.

It is essential to acknowledge that any observed variation might be due to differences in methodologies at different scales, but this likely introduces noise rather than bias.

### 2.2 | Collating Published Camera Studies

We compiled published camera trap records by searching Web of Science with the following criteria: ‘camera\*’ AND Asia\* or Thai\* or Malaysia\* or Indonesia\* or Borneo\* or Cambodia\* or Vietnam\* or Lao\* or Myanmar\* or Burm\* or Sumatra\*. We selected from the list of returned studies those that were written in English and reported relevant results for the sun bear, including sampling effort (number of cameras, and deployment length or total trap nights), and number of independent detections using a 30–60 min interval between detections of the same species, referred to as “independence period” (Rovero and Zimmermann 2016). We examined the references listed in key papers to identify and include further sources. We included all tropical forest camera studies that used unbaited cameras placed < 0.4 m high, usually facing trails or other areas used by wildlife. This is the standard deployment approach used in the region and suitable for most semi-terrestrial species > 1 kg (Rovero and Zimmermann 2016). From each study, we recorded the location (forest name and coordinates), detection, and effort (number of trap nights). We grouped multiple studies from the same landscape in the same year by summing detections and effort among the studies and averaging the covariate values (Greco et al. 2025). We assumed that all studies that occurred in regions where the sun bear is sympatric with the Asiatic black bear (*Ursus thibetanus*) correctly distinguished between these two species. The full dataset is publicly available (Mendes et al. 2024a).

### 2.3 | New Camera Trapping Sessions

We conducted a total 19 new camera trapping sessions in eight tropical forest landscapes in Thailand, Peninsular Malaysia, Sumatra, and Borneo between December 2013 and June 2020. These sessions occurred in seven different landscapes in the sun bear's range as recognised by the IUCN (Scotson et al. 2017a). One session of camera trapping also occurred outside the sun bear's IUCN range, in Lambir Hills National Park in Sarawak (taking the total number of landscapes where new camera trapping occurred to eight). Further details on the new camera trapping sessions are provided in Supporting Information: Table S1.

We deployed between 18 and 78 passive infrared Bushnell Trophy and Reconyx HC500 Hyperfire cameras across sampling areas ranging from 10 to 813 km<sup>2</sup>. We standardised deployment methods across all landscapes. Cameras were placed within a pre-mapped grid and spaced at least 500 m apart in large, forested landscapes (> 50 km<sup>2</sup>) and 100–500 m apart in smaller forest patches. Cameras were attached to trees 0.3 m above ground along hiking trails or natural wildlife trails and

deployed for approximately 60–90 days. Sun bear detections were considered independent if they occurred at least 30 min apart (Rovero and Zimmermann 2016). We differentiated sun bears from Asiatic black bears based on body shape and size, as well as fur colour. Research permissions are provided in Supporting Information: Table S7.

## 2.4 | Landscape-Level Habitat Associations Using GLMMs of Sun Bear Detections

We used both the published and new camera trap data to investigate landscape-scale relationships between the number of sun bear detections and various environmental and anthropogenic factors using generalised linear mixed-effect models (GLMMs) with a negative binomial error distribution (Mendes and Luskin 2025). We treated detections as count data and included a fixed continuous term to control for study effort (measured in trap nights) and a random categorical term for landscape, because some landscapes had multiple studies. Following Ash et al. (2020), we used raw count data as opposed to a relative abundance index (RAI, independent photos per 100 trap nights) to enable use of a negative binomial distribution. We note that these approaches do not account for variation in detection probability, and thus we are implicitly assuming that detection probability among studies does not vary, and acknowledge this may introduce measurement error and that detections do not linearly reflect true abundance (Sollmann et al. 2013). We also acknowledge that there is unexplained variation in detections owing to slight differences among studies in equipment, deployments, and data curation. Furthermore, sun bear detectability could vary due to the availability of fruit across time, such as in masting years when there may be a reduced need to travel (Fredriksson et al. 2006; Dehaudt et al. 2024b). All sources of measurement error reduce our power of detecting statistically significant or ‘true’ relationships, but do not lend themselves to directional bias.

We used GLMMs to test the effect of 10 environmental and anthropogenic predictor variables (covariates) on sun bear detections reported in published camera studies conducted across different landscapes. Our covariate values were derived from GIS layers and describe the area within a 20 km radius around the centroid of each landscape. Forest sizes were determined from the IUCN Red List protected areas database (UNEP-WCMC and IUCN 2021) or, for unprotected areas, from the original papers’ reporting. Sources for covariate data are listed in Supporting Information: Table S2. We used this study area (1256 km<sup>2</sup>) to account for large sampling grids and unclear centroids inferred for some landscapes. We also used the year of camera trapping as a covariate (resulting in 11 total covariates tested) to determine if and how the amount of sun bear detections has changed over time. Details of the references, study sites, camera effort, and sampling periods are available in the CamTrapAsia database (Mendes et al. 2024a).

We tested each variable independently with linear and non-linear (quadratic) univariate models and used AICc model selection to identify the most parsimonious models (Burnham and Anderson 2002). We used the conventional cut-off of 2.0 to identify clear differences in models. Nonlinear models allowed

us to identify peaks of sun bear detections that could exist at mid-levels of a covariate value. We also tested bivariate models. To avoid multicollinearity, we tested correlation among our covariate pairs by creating a correlation matrix and excluding all covariate pairs that had a correlation coefficient of 0.4 or greater. All GLMMs were implemented using the R package “glmmTMB” (version 1.1.8) (Brooks et al. 2017).

## 2.5 | Local-Scale Habitat Associations Using Royle–Nichols Hierarchical Modelling

We assessed the effect of habitat variables on sun bear abundance at the local scale using Royle–Nichols abundance models (Royle and Nichols 2003) (see Supporting Information: Table S4 for details of sampling effort, detection rates, and naïve occupancy). For this, we only used data from our 19 new camera trapping surveys. We use the Royle–Nichols model, which uses presence–absence data to derive a relative abundance metric by exploiting the positive relationship between variation in individual detection probability and the species’ abundance (Royle and Nichols 2003). RN models reveal habitat associations by estimating abundance relative to a gradient of a covariate while accounting for imperfect detection (Kéry and Royle 2021). Hierarchical abundance modelling, such as N-mixture models, often face assumption violations when using camera trap data, so we conservatively interpret the results of our model and avoid inferring that these values reflect the actual abundance or density. Rather, relative abundance is only informative when being predicted by covariates to reflect spatial variation in abundance (Gilbert et al. 2021).

To ensure that model outputs were spatially comparable across multiple landscapes and to prevent detecting the same sun bear at different sampling units, we resampled the data into 10 km<sup>2</sup> hexagon grid cells, as suggested by Rayan and Linkie (2020), which is smaller than one estimate of the average home range of sun bears of 14.8 km<sup>2</sup> (Te Wong et al. 2004). We produced a sun bear detection history matrix based on a sampling occasion of 3 days containing presence/absence data (0 = sun bear not detected; 1 = sun bear detected; NA = inactive sampling unit or occasion). Each sampling unit (hexagon) contained at least one camera, but for those with two or more, sun bear presence/absence per sampling unit and sampling occasion was informed by all active cameras in the sampling unit they occurred in, and we averaged their covariate values. We included “study” as a fixed effect to account for potential variation in abundance associated with each set of deployments, and to maintain the spatial and temporal independence of our sampling units, as well as satisfy the assumption of population closure. We also included the total effort, in trap nights, per sampling unit as a fixed effect in the detection formula to account for multiple cameras grouped in a spatially explicit sampling unit (e.g., multiple cameras in one grid cell). We tested the effect of local-scale predictors such as elevation, percentage of oil palm plantations, human population, and percentage of degraded forest, which were all calculated using a 1 km radius around each camera’s coordinates (3.14 km<sup>2</sup> area; sources for all covariate data are listed in Supporting Information: Table S2). Degraded forest included ‘combined land cover of oil palm, lowland mosaics, lowland open ground, and regrowth/

plantations' and is sourced from Miettinen et al. (2016). The variable present in the GLMM analysis but not the RN analysis were forest size, forest cover, roughness, and night lights. We addressed potential issues with collinearity among covariates by limiting our investigations to univariate models as we did not obtain sufficient detections at many sites to test multivariate models. We used AICc to identify the most parsimonious univariate model and implemented all analyses in the R packaged 'unmarked' (Fiske and Chandler 2011). The conventional cut off of 2.0 AICc points was again used to identify clear differences in models.

## 2.6 | Diel Activity Patterns

We used time-stamped detections from our new sampling to investigate if forest degradation affects the sun bear's diel activity by comparing Kernel density estimates of sun bears in degraded and non-degraded (intact) forests. Specifically, we split detections based on the median value of the forest degradation variable (sourced from Miettinen et al. 2016) and ran a bootstrap procedure to simulate 1000 distributions of activity pattern data to conduct a Wald test using the function compareAct() in the R package "activity" (Rowcliffe et al. 2014). For this analysis, we calculated forest degradation at a 1.0 km radius around each camera (3.14 km<sup>2</sup>) because diel activity is more likely to be affected by local conditions. The coefficient of overlap was calculated using the R package 'overlap' (Ridout and Linkie 2009).

## 3 | Results


### 3.1 | Landscape-Level Habitat Associations

From both the new and previously published camera trap studies, we obtained 3360 independent detections of sun bears from 127 camera studies at 38 landscapes in six different countries, as well as 58 studies in the sun bear's range that did not detect the species (Table 1, Figure 1). The bivariate GLMM model containing forest size and night lights (a measure of proximity to humans) was the best predictor of sun bear detections (Table 2 and Supporting Information: S5), with a significant positive relationship for both covariate terms ( $p < 0.001$  for forest size and  $p = 0.014$  for night lights; Figure 2, Table 2). Examples of forests in the top 20% of sun bear detection rate, human population, and night lights were in Sumatra (Bukit Barisan Selatan National Park, the eastern area of the Gunung Leuser Ecoregion, and Kerinci Seblat National Park) and central Peninsular Malaysia (eastern area of Taman Negara and near Jerneh). Similar to the night lights

variable, the human population also had a significant positive influence on sun bear detections ( $p = 0.03$ , Table 2 and Supporting Information: S5).

### 3.2 | Local-Scale Habitat Associations

There were 416 independent detections of sun bears in our 19 new camera trapping sessions across eight landscapes, including detections of multiple sun bears together (usually mothers with a single cub). Using the Royle–Nichols (RN) abundance models, elevation was the best predictor (2.88 AICc points lower than the next best performing model) and produced a significant positive relationship ( $p < 0.001$ , Table 3, and Supporting Information: S6; Figure 3a). The next best models included a



**FIGURE 1** | The sun bear's historic range (shown in orange) with points showing the location of sun bear occurrence records used in this study. The historic range data is sourced from the IUCN Red List (Scotson et al. 2017a).

**TABLE 1** | Data sources and sample sizes (detections) for the three analyses of this study. Data from new camera trapping data sets were also included in the GLMM analyses.


| Analysis   | Records     |
|--|-------------|
| Camera trap count data for GLMMs (landscapes, surveys with detections, surveys without detections) | 38, 127, 58 |
| Independent detections for GLMMs   | 3360        |
| New camera data for RN models and diel activity (landscapes, surveys)                              | 8, 19       |
| Independent detections for RN models and diel activity   | 416         |

Abbreviations: GLMMs, generalised linear mixed-effect models; RN, Royle and Nichols.

**TABLE 2** | Top five GLMM models assessing sun bear detections among landscapes as determined by AICc model selection.

| Model ranking | Terms                  | Estimate | Std. error | p-value |
|---------------|------------------------|----------|------------|---------|
| Model 1       | Forest size (log)      | 0.50     | 0.11       | < 0.01* |
|               | Night lights           | 0.38     | 0.16       | 0.01*   |
| Model 2       | Forest size (log)      | 0.49     | 0.11       | < 0.01* |
|               | Human population (log) | 0.29     | 0.14       | 0.03*   |
| Model 3       | Forest size (log)      | 0.42     | 0.11       | < 0.01* |
|               | Year                   | -0.31    | 0.15       | 0.04*   |
| Model 4       | Forest size (log)      | 0.44     | 0.13       | < 0.01* |
|               | Elevation              | -0.06    | 0.14       | 0.64    |
| Model 5       | Forest size (log)      | 0.39     | 0.12       | < 0.01* |
|               | Forest cover           | 0.06     | 0.17       | 0.74    |

Note: We used AICc model selection criteria to compare negative binomial GLMMs. The full model ranking table, which includes the AICc and  $\Delta$ AICc values, is Supporting Information: Table S5. The response variable was independent sun bear detections in 185 published camera trap studies, and we included study effort as a fixed effect and landscape as a random effect. All covariates were averaged for the 20 km radius area surrounding the study site, then centred and standardised so effect sizes could be interpreted relative to each other, with the exceptions of Forest size, which was measured in km<sup>2</sup> and year, which refers to the year that a camera trapping study began. Correlations among predictors were checked and predictor pairs that had a correlation coefficient of 0.4 or greater were included. Only linear covariates were used for the multivariate models. Sources for covariate data are listed in Supporting Information: Table S2. The asterisks (\*) denotes significant results. Abbreviations: AIC, Akaike information criterion; GLMMs, generalised linear mixed-effect models.



**FIGURE 2** | Sun bear detections (per 100 trap nights) from 185 published and new camera trap studies, as predicted by the top GLMM (Table 2), which contained both forest size and night lights as predictors. The blue line shows the model at the 95th percentile night light value associated with a camera trap survey conducted in the sun bear's range, whereas the red line shows the model at the 5th percentile for night lights. The blue and red ribbons represent the 95% confidence interval for both conditions, however there is considerable overlap between these ribbons resulting in a purple ribbon and only a small section of red ribbon in the bottom right corner of the graph. GLMM, generalised linear mixed-effect model.

positive effect of human population ( $\Delta$ AICc = 2.88;  $p = 0.001$ ) and a negative effect of oil palm ( $\Delta$ AICc = 5.51;  $p = 0.024$ ; Table 3, and Supporting Information: S6; Figure 3b,c). Intact and degraded forests had a positive and negative influence on sun bear abundance, respectively, although these were not the best performing models based on AICc model selection (Supporting Information: Table S6).

**TABLE 3** | Top five abundance models assessing sun bear habitat associations at a local scale (within landscapes), as determined by AICc model selection.

| Terms             | Estimate | Std. error | p-value |
|-------------------|----------|------------|---------|
| Elevation         | 0.3      | 0.09       | < 0.01* |
| Human population  | 0.25     | 0.08       | < 0.01* |
| Oil palm          | -0.31    | 0.14       | 0.02*   |
| Forest intactness | 0.32     | 0.17       | 0.07    |
| Degraded forest   | -0.28    | 0.17       | 0.1     |


Note: The full model ranking table, which includes the AICc and  $\Delta$ AICc values, is Supporting Information: Table S6. We fit Royle-Nichols abundance models with camera trapping session as a covariate to account for variation among landscapes and between surveys in the same landscape and effort affecting detection probability (per 10 km<sup>2</sup> sampling unit). Sources for covariate data are listed in Supporting Information: Table S2. The asterisks (\*) denote significant results. Abbreviation: AIC, Akaike information criterion.

### 3.3 | Diel Activity Patterns


Using the 416 independent detections of sun bears from our new camera trapping, we assessed sun bear diel activity using kernel density estimates. Sun bears showed a clear bi-modal activity peaks inferring a crepuscular activity pattern, with a first activity peak at dawn and a more prominent activity peak at dusk and shortly thereafter (Figure 4). For our activity pattern comparison between sun bears living in degraded and non-degraded forests, the median forest degradation value was zero. There were 724 cameras classified as intact forest and 372 cameras classified as degraded (noting that we excluded degraded landscapes with no sun bear detections). There was no change in diel activity when comparing intact versus degraded forests (Wald test  $p = 0.271$ ; Figure 4).

## 4 | Discussion

Using detections sourced from 127 camera trap studies from 38 landscapes, sun bear detections were highest in large forests



**FIGURE 3** | Local variation in sun bear predicted abundance determined from detection-correction Royle–Nichols hierarchical modelling of camera trapping from eight landscapes in Southeast Asia (the three best performing models are shown; Table 3 and Supporting Information: S6). The y-axis is predicted sun bear relative abundance per 10 km<sup>2</sup> sampling unit. All covariates were centred and standardised prior to modelling, so effect sizes can be compared between variables.



**FIGURE 4** | Diel activity pattern of the sun bear in degraded (orange) and non-degraded forests (blue) determined from 416 independent detections during 19 camera trapping surveys at eight landscapes across their range in Southeast Asia. Time-stamped sun bear detections were assigned to the degraded or non-degraded forest category based on whether the detection occurred in intact versus degraded forest. Degradation included any forest with oil palm plantations, lowland mosaic forest, lowland open ground, and regrowth forests within a 1.0 km radius of a camera trap in our new camera trapping sessions (Supporting Information: Table S2).  $\Delta_1$  refers to the extent of overlap in activity of sun bears in the two different forest categories.  $W$  refers to the Wald test result comparing these two distributions and  $p$  refers to the  $p$  value associated with this Wald test. The tick marks (rugs) at the bottom of the figure show the exact time of each sun bear detection, coloured-coded to match the degradation status of the location in which it occurred.

and areas with more night lights, a proxy for humans and their infrastructure. Two notable areas with high detections of sun bears near night lights (humans) were landscapes in Sumatra and central Peninsular Malaysia, where there are many towns directly bordering large parks. This result was consistent with the local scale Royle–Nichols (RN) hierarchical modelling from eight landscapes, which also showed sun bears being positively associated with humans and also elevation, but negatively associated with oil palm plantations. Taken together, a parsimonious explanation for these results is that sun bears are

frequently detected at the edges of large forests in areas close to humans (where there are night lights), and within these large forested landscapes (local scale), they show a preference for higher elevations and an avoidance of oil palm plantations. These results provide a rationale to be cautiously optimistic for the conservation of the species in degraded forests.

A key inference from this work was that forest size appeared more important for sun bears than forest quality. The importance of large forests may be due to their large 15 km<sup>2</sup> home ranges and low densities (Te Wong et al. 2004; Abidin et al. 2018). Hwang et al. (2021) also proposed that sun bears can persist in disturbed forests because the greater amount of dead wood in such forests supports insects for sun bears to feed on (Luke et al. 2014; Kimber and Eggleton 2018), and invertebrates comprise a major part of the sun bear diet, particularly in non-masting years when there is less fruit (Wong 2002; Te Wong et al. 2005).

The positive associations with humans in both analyses were unexpected because sun bears are poached, and most megafauna populations decline near humans (Shepherd and Shepherd 2010; Benítez-López et al. 2017; Hughes 2017; Ripple et al. 2019; Gomez et al. 2020; Edinur et al. 2022). The positive associations with humans may be an artifact of high detection of sun bears in Sumatra and Peninsular Malaysia, where there happened to be many towns bordering the large parks. If true, sun bears were not “attracted” to areas with more people and lights, but may be able to survive there.

The negative influence of oil palm was expected, but still interesting because sun bears are known to forage in oil palm plantations (Normua et al. 2004; Abidin et al. 2018; Guharajan et al. 2018). This is likely due to the increased risk of snaring and the access for poachers that oil palm plantations provide and we note that sun bears that utilise plantations still require nearby forests (Cheah 2013; Azhar et al. 2014; Guharajan et al. 2018).

Sun bears are primarily frugivores, and megafauna seed dispersers are crucial for maintaining forest tree community dynamics (Dehadt et al. 2024a; Anderson et al. 2025; Deblauwe et al. 2025). Large frugivores are often at higher risk (Joshi et al. 2025) and there are numerous ecological cascades that may result from declines in large seed dispersers (Luskin et al. 2017; Williams et al. 2021; Brodie et al. 2025).

## 4.1 | Scale-Dependency

We expected there to be variation in the strength of habitat covariates among our analyses because sun bear local habitat preferences may differ from those of the entire region they occupy, and the analyses used different data. Interpreting multiple analyses together can provide richer insights but requires nuance. A key example was elevation, which was not a key variable in the landscape-level analyses (Supporting Information: Table S5) but was the top variable in the local-scale analysis (Supporting Information: Table S6). One explanation is that the local analysis cameras were predominantly placed between 0 and 800 m asl, while the landscape-level studies had a range of 9 to 5000 m asl. Taken together, this may indicate a preference for hill forests (e.g., 400–800 m asl) and an avoidance of the lowest and highest areas of the sun bear's range, such as peat swamps < 50 m asl and montane forests > 1000 m asl. This finding is consistent with other studies (Linkie et al. 2007; Steinmetz et al. 2011, Wong et al. 2013).

## 4.2 | Sun Bear Diel Activity

Many animals—especially persecuted megafauna—adopt a more nocturnal activity pattern in degraded areas to avoid encounters with humans (Gaynor et al. 2018; Lee et al. 2024). For example, both Tee et al. (2021) and Fredriksson (2012) found that sun bears living near forest edges are more nocturnal than conspecifics residing in forest interiors. Instead, we found sun bears were consistently crepuscular (Figure 4). This is consistent with Guharajan et al. (2018) but conflicts with Te Wong et al. (2004) and Cheah (2013), who reported sun bears as being diurnal but become more nocturnal when feeding in oil palm plantations. Methodological differences may explain these inconsistent results, as Guharajan et al. (2018) and Tee et al. (2021) utilised camera traps while Fredriksson (2012) and Te Wong et al. (2004) used radio collars. Future studies utilising radio or GPS collars could resolve discrepancies in sun bear behaviour near humans and forest degradation.

## 4.3 | Comparison to Previous Studies

Aside from Scotson et al. (2017b), previous analyses of sun bear detection data have been limited to small geographic areas (Guharajan et al. 2021; Hwang et al. 2021; Guharajan et al. 2022). These studies identified that canopy cover and other comparable habitat covariates are positively associated with sun bear detections and persistence in moderately degraded habitat (Hwang et al. 2021). Our study expands on these previous studies by sourcing sun bear camera trap detections from multiple countries and also analysing more covariates than these previous studies, including both habitat and anthropogenic covariates. A key covariate that was analysed in this study was the extent of oil palm plantations. The conversion of rainforest into oil palm plantations is a major conservation issue in the region, so understanding the response of sun bears to oil palm plantations is crucial for their conservation. Overall, our paper concurs with Scotson et al. (2017b) and Guharajan et al. (2022) who identified that large expanses of forested habitat are

important for sun bears, but the species can also persist in areas with moderate degradation (Guharajan et al. 2021; Hwang et al. 2021). Together, these findings suggest that sun bears may not avoid humans in space or time.

## 4.4 | Sun Bear Conservation

Scotson et al. (2017b) and Wong et al. (2013) suggest recent sun bear declines are due to forest loss and poaching (Edinur et al. 2022). Approximately 90% of the potentially suitable habitat for sun bears lies outside protected areas, where poaching may occur (Scotson 2019). Our analyses suggest there is a considerable opportunity to safeguard the species by targeting the protection of large, unprotected forests, which is consistent with other recent analyses in the region (Scotson et al. 2017b; Hwang et al. 2021). Namely, there are large expanses of natural and native timber production forests in the region that may be targeted for this purpose, alongside the enforcement of regulations against poaching and wildlife trade (Linkie et al. 2015; Edinur et al. 2022).

One ray of hope comes from Lambir Hills National Park in Sarawak, Malaysian Borneo, which contains 70 km<sup>2</sup> of intact lowland and hill forest that previously experienced severe poaching. Sun bears were not detected in camera surveys conducted at Lambir Hills in 2002 and 2006, suggesting they may have been extirpated (Shahanan and Debski 2002; Azlan and Lading 2006). There has since been a renewed emphasis on park management and enforcement to reduce poaching, and our 2017 survey detected a sun bear at Lambir Hills National Park. Another potential positive is that the recent outbreak of African Swine Fever may reduce competition with wild boars and improve foraging opportunities for sun bears (Luskin et al. 2021; Luskin et al. 2023b; Lieb et al. 2025).

## 4.5 | Future Research Directions

To determine whether degraded forests are population sinks or sustain viable populations, longitudinal work across the degradation gradient is needed (Dehaut et al. 2025). The continued use of cameras among many different groups presents opportunities for collaborations to achieve longitudinal work (Bruce et al. 2025). We also recommend that more camera trap surveys be conducted in regions north of Peninsular Malaysia (e.g., Thailand, Myanmar, and Cambodia), as much of the data for this study was sourced from Peninsular Malaysia, Sumatra, and Borneo. Movement studies to understand how sun bears use degraded forests and corridors will enable better conservation planning. Diet studies are also required to test the hypothesis that sun bears become more insectivorous in degraded forests (Hwang et al. 2021) and the degree to which they consume oil palm and other agricultural products in degraded forests (Cheah 2013). More information on sun bear health, such as variation in their body condition and reproduction, would be especially useful, particularly if such research spans masting and non-masting years and is repeated in intact versus degraded forests. Finally, there is a need to determine whether sun bears are habituating to humans at forest edges and whether human-wildlife conflict is occurring, such as through altered diets.

## Author Contributions

**Alexander Hendry:** formal analysis, writing – original draft preparation, writing – review and editing. **Harsh K. Pahuja:** formal analysis, writing – original draft preparation, writing – review and editing. **Zachary Amir:** data curation, formal analysis, methodology, supervision, visualisation, writing – review and editing. **Ilyas Nursamsi:** formal analysis, visualisation. **Jonathan H. Moore:** data curation, formal analysis, fieldwork. **Matthew Scott Luskin:** conceptualisation, data curation, formal analysis, funding acquisition, fieldwork, methodology, supervision, writing – original draft preparation, writing – review and editing.

## Acknowledgements

The research was funded by the Smithsonian Institution's ForestGEO programme, Nanyang Technological University, the University of Queensland, and the National Geographic Society Committee for the Research and Exploration #9384–13. In kind support was provided by Fauna and Flora International's Indonesia programme, Institution Conservation Society (ICS)—Solok Selatan Wahana Konservasi Masyarakat, the Leuser International Foundation (LIF). We thank Yayasan Sabah, the Sabah Forest Department, the Sabah Biodiversity Council, and the Danum Valley Management Committee, Abdul Hamid, Glen Reynolds, Jedediah Brodie, Katie Doehla, and Tombi Karolus for permission and help conducting fieldwork at Danum Valley. We thank Patrick Jansen, Jorge Ahumada, the Smithsonian Institution, the Tropical Ecology Assessment and Monitoring (TEAM) network, Yao Tse Leong and the Forest Research Institute Malaysia (FRIM) for permissions and data collection at Pasoh. We thank Mohizah Bt. Mohamad, Januarie Kulis and the Sarawak Forestry Department and the NTU field ecology courses for permission and help with fieldwork data at Lambir Hills. We thank Shawn Lum, Adrian Loo, Max Khoo, Ben Lee, Jasyln Chan, Alexis Goh, and NParks for permission and help with fieldwork in Singapore. We thank Wirong Chantorn, Anuttara Nathalang, Sarayudh Bunyavejchewin, Ronglarp Sukmasuang, Feliz Gutierrez, Chris Scanlon for permissions and help at Khao Yai and Khao Ban Tat/Khao Chong. We thank Wido Rizqi Albert, Matthew Linkie, Yoan Dinata, Hariyo Wibisono and HarimauKita for help facilitating fieldwork in Sumatra, and we thank Edi Siarenta Sembiring, Tarmizi and Eka Ramadyanta, Salpayanri, Iswandri Tanjung and Chris Decky for assistance with fieldwork. We thank the members of the Ecological Cascades Lab and anonymous reviewers for comments that improved previous drafts.

## Ethics Statement

Ethical approval for non-invasive camera sampling was not required for this study.

## Conflicts of Interest

The authors declare conflicts of interests.

## Data Availability Statement

Data available in FigShare: [https://figshare.com/articles/dataset/Sun\\_Bear\\_Data/23993505](https://figshare.com/articles/dataset/Sun_Bear_Data/23993505). The data that support the findings of this study are openly available in FigShare at [https://figshare.com/articles/dataset/Sun\\_Bear\\_Data/23993505](https://figshare.com/articles/dataset/Sun_Bear_Data/23993505), reference number 23993505.

## References

Abidin, M. K. Z., A. A. Mohammed, and S. M. Nor. 2018. AIP Conference Proceedings.

Amir, Z., J. H. Moore, P. J. Negret, and M. S. Luskin. 2022. "Megafauna Extinctions Produce Idiosyncratic Anthropocene Assemblages." *Science Advances* 8: eabq2307.

Anderson, S. E., Z. Amir, T. Bruce, and M. S. Luskin. 2025. "Range-Wide Camera Trapping for the Australian Cassowary Reveals Habitat

Associations With Rainfall and Forest Quality." *Ecology and Evolution* 15: e71464.

Ash, E., Ž. Kaszta, A. Noochdumrong, T. Redford, and D. W. Macdonald. 2020. "Environmental Factors, Human Presence and Prey Interact to Explain Patterns of Tiger Presence in Eastern Thailand." *Animal Conservation* 24: 12631.

Augeri, D. M. 2005. *On the biogeographic ecology of the Malayan sun bear*. University of Cambridge Cambridge.

Azhar, B., D. B. Lindenmayer, J. Wood, J. Fischer, and M. Zakaria. 2014. "Ecological Impacts of Oil Palm Agriculture on Forest Mammals in Plantation Estates and Smallholdings." *Biodiversity and conservation* 23: 1175–1191.

Azlan, M. J., and E. Lading. 2006. "Camera Trapping and Conservation in Lambir National Park, Sarawak." *Raffles Bulletin of Zoology* 54, no. 2: 469–475.

Benítez-López, A., R. Alkemade, A. M. Schipper, et al. 2017. "The Impact of Hunting on Tropical Mammal and Bird Populations." *Science* 356: 180–183.

Brodie, J. F., C. Bello, C. Emer, et al. 2025. "Defaunation Impacts on the Carbon Balance of Tropical Forests." *Conservation Biology* 39: e14414.

Brodie, J. F., A. J. Giordano, E. F. Zipkin, H. Bernard, J. Mohd-Azlan, and L. Ambu. 2015. "Correlation and Persistence of Hunting and Logging Impacts on Tropical Rainforest Mammals." *Conservation Biology* 29: 110–121.

Brodie, J. F., J. Mohd-Azlan, C. Chen, et al. 2023. "Landscape-Scale Benefits of Protected Areas for Tropical Biodiversity." *Nature* 620: 807–812.

Brooks, E., K. Kristensen, J. v Benthem, et al. 2017. "Glmmtmb Balances Speed and Flexibility Among Packages for Zero-Inflated Generalized Linear Mixed Modeling." *R Journal* 9: 378.

Bruce, T., Z. Amir, B. L. Allen, et al. 2025. "Large-Scale and Long-Term Wildlife Research and Monitoring Using Camera Traps: A Continental Synthesis." *Biological Reviews* 100: 530–555.

Burnham, K. P., and D. R. Anderson. 2002. *Model Selection and Multimodel Inference: A Practical Information-theoretic Approach* (Second edition. edition). Springer.

Carr, E., Z. Amir, C. P. Mendes, J. H. Moore, I. Nursamsi, and M. S. Luskin. 2023. "The Highs and Lows of Serow (*Capricornis sumatraensis*): Multi-Scale Habitat Associations Inform Large Mammal Conservation Strategies in the Face of Synergistic Threats of Deforestation, Hunting, and Climate Change." *Raffles Bulletin of Zoology* 71: 400–416.

Cheah, C. 2013. *The Ecology of Malayan Sun Bears at the Krau Wildlife Reserve, Pahang, Malaysia and Adjacent Plantations*. PhD dissertation, University Putra Malaysia.

Clements, G. R., A. J. Lynam, D. Gaveau, et al. 2014. "Where and How Are Roads Endangering Mammals in Southeast Asia's Forests?" *PLoS One* 9: e115376.

Deblauwe, V., M. S. Luskin, S. D. Assola, et al. 2025. "Declines of Ebony and Ivory Are Inextricably Linked in An African Rainforest." *Science Advances* 11: eady4392.

Decœur, H., Z. Amir, C. P. Mendes, J. H. Moore, and M. S. Luskin. 2023. "Mid-Sized Felids Threatened by Habitat Degradation in Southeast Asia." *Biological Conservation* 283: 110103.

Dehault, B., Z. Amir, and H. Decoeur, et al. 2022. "Common Palm Civets *Paradoxurus Hermaphroditus* Are Positively Associated With Humans and Forest Degradation With Implications for Seed Dispersal and Zoonotic Diseases." *Journal of Animal Ecology* 91: 794–804.

Dehault, B., R. Bose, J. J. Avoto, et al. 2025. "Thirty Years of Arboreal Wildlife Trends in An African Rainforest Under Evolving Threats and Researchers' Presence." *Biological Conservation* 312: 111475.

- Dehaudt, B., T. Bruce, V. Deblauwe, et al. 2024a. "Divergent Seed Dispersal Outcomes: Interactions Between Seed, Disperser, and Forest Traits." *Ecology* 105: e4409.
- Dehaudt, B., J. Moore, L. Gibson, and M. S. Luskin. 2024b. "More Problems Sampling Wildlife With Baits: Environmental Food Resources Influence Trapping Rates." *Wildlife Letters* 2: 83–89.
- Dunn, A., Z. Amir, H. Decoeur, et al. 2022. "The Ecology of the Banded Civet (*Hemigalus derbyanus*) in Southeast Asia With Implications for Mesopredator Release, Zoonotic Diseases, and Conservation." *Ecology and Evolution* 12: e8852.
- Edinur, H. A., C. G. N. Hajar, and M. T. Abdullah. 2022. "COVID-19 Treatment May Impact Sun Bear." *Journal of Sustainability Science and Management* 17: 1–5.
- Fiske, I., and R. Chandler. 2011. "Unmarked": An R Package for Fitting Hierarchical Models of Wildlife Occurrence and Abundance." *Journal of Statistical Software* 43: 1–16.
- Fotang, C., U. Bröring, C. Roos, et al. 2021. "Human Activity and Forest Degradation Threaten Populations of the Nigeria–Cameroon Chimpanzee (*Pan Troglodytes Elliotti*) in Western Cameroon." *International Journal of Primatology* 42: 105–129.
- Fredriksson, G. M. 2012. *Effects of El Niño and large-scale forest fires on the ecology and conservation of Malayan sun bears (Helarctos malayanus) in East Kalimantan, Indonesian Borneo*. Universiteit van Amsterdam [Host].
- Fredriksson, G. M., S. A. Wich, and Trisno. 2006. "Frugivory in Sun Bears (*Helarctos malayanus*) Is Linked to El Niño-Related Fluctuations in Fruiting Phenology, East Kalimantan, Indonesia: Phenological Influences on Sun Bear Frugivory." *Biological Journal of the Linnean Society* 89: 489–508.
- Gallego-Zamorano, J., A. Benítez-López, L. Santini, J. P. Hilbers, M. A. J. Huijbregts, and A. M. Schipper. 2020. "Combined Effects of Land Use and Hunting on Distributions of Tropical Mammals." *Conservation Biology* 34: 1271–1280.
- Gaynor, K. M., C. E. Hohnowski, N. H. Carter, and J. S. Brashares. 2018. "The Influence of Human Disturbance on Wildlife Nocturnality." *Science* 360: 1232–1235.
- Gilbert, N. A., J. D. J. Clare, J. L. Stenglein, and B. Zuckerberg. 2021. "Abundance Estimation of Unmarked Animals Based on Camera-Trap Data." *Conservation Biology* 35: 88–100.
- Gomez, L., C. Shepherd, and M. Khoo. 2020. "Illegal Trade of Sun Bear Parts in the Malaysian States of Sabah and Sarawak." *Endangered Species Research* 41: 279–287.
- Gouda, S., N. S. Chauhan, J. Sathy, and H. Sahu. 2020. "Daily Activity Pattern of Malayan Sun Bear in Dampa Tiger Reserve, Mizoram, India." *Journal of Wildlife and Biodiversity* 4: 56–64.
- Grantham, H. S., A. Duncan, T. D. Evans, et al. 2020. "Anthropogenic Modification of Forests Means Only 40% of Remaining Forests Have High Ecosystem Integrity." *Nature Communications* 11, no. 1: 5978. <https://doi.org/10.1038/s41467-020-19493-3>.
- Greco, I., L. Beaudrot, C. Sutherland, et al. 2025. "Landscape-Level Human Disturbance Results in Loss and Contraction of Mammalian Populations in Tropical Forests." *PLoS Biology* 23: e3002976.
- Griffiths, M., and C. P. Schaik. 1993. "The Impact of Human Traffic on the Abundance and Activity Periods of Sumatran Rain Forest Wildlife." *Conservation Biology* 7: 623–626.
- Guharajan, R. 2016. *Survival Strategies of the Sun Bear (Helarctos malayanus) in the Lower Kinabatangan Floodplain, Sabah, Malaysian Borneo*. University of Minnesota.
- Guharajan, R., J. Abrams, and N. Abram, et al. 2022. "Determinants of Sun Bear *Helarctos malayanus* Habitat Use in Sabah, Malaysian Borneo and Its Predicted Distribution under Future Forest Degradation and Loss." *Biodiversity and conservation* 32: 02503-9.
- Guharajan, R., T. W. Arnold, G. Bolongon, et al. 2018. "Survival Strategies of a Frugivore, the Sun Bear, in a Forest-Oil Palm Landscape." *Biodiversity and Conservation* 27: 3657–3677.
- Guharajan, R., A. Mohamed, S. T. Wong, et al. 2021. "Sustainable Forest Management Is Vital for the Persistence of Sun Bear *Helarctos Malayanus* Populations in Sabah, Malaysian Borneo." *Forest Ecology and Management* 493: 119270.
- Haddad, N. M., L. A. Brudvig, J. Clobert, et al. 2015. "Habitat Fragmentation and Its Lasting Impact on Earth's Ecosystems." *Science Advances* 1, no. 2). <https://doi.org/10.1126/sciadv.1500052>.
- Harrison, R. D., R. Sreekar, J. F. Brodie, et al. 2016. "Impacts of Hunting on Tropical Forests in Southeast Asia." *Conservation Biology* 30: 972–981.
- Hendry, A., Z. Amir, and H. Decoeur, et al. 2023. "Marbled Cats in Southeast Asia: Are Diurnal and Semi-Arboreal Felids at Greater Risk From Human Disturbances?" *Ecosphere* 14: e4338.
- Honda, A., Z. Amir, C. P. Mendes, J. H. Moore, and M. S. Luskin. 2023. "Binturong Ecology and Conservation in Pristine, Fragmented and Degraded Tropical Forests." *Oryx* 1: 1–10.
- Hughes, A. C. 2017. "Understanding the Drivers of Southeast Asian Biodiversity Loss." *Ecosphere* 8: e01624.
- Hwang, M. H., M. A. Ditmer, S. D. Teo, S. T. Wong, and D. L. Garshelis. 2021. "Sun Bears Use 14-Year-Old Previously Logged Forest More Than Primary Forest in Sabah, Malaysia." *Ecosphere* 12: e03769.
- Joshi, P. R., A. Semper-Pascual, A. F. Vallejo-Vargas, et al. 2025. "Species Characteristics Modulate Tropical Mammal Responses to Forest Quality in Southeast Asia." *Biological Conservation* 310: 111388.
- Kéry, M., and J. A. Royle. 2021. *Applied Hierarchical Modeling in Ecology Modeling Distribution, Abundance and Species Richness in R and BUGS* (1st edition). Elsevier Academic Press.
- Kimber, A., and P. Eggleton. 2018. "Strong but Taxon-Specific Responses of Termites and Wood-Nesting Ants to Forest Regeneration in Borneo." *Biotropica* 50: 266–273.
- Krishnasamy, K., and C. R. Shepherd. 2014. "A Review of the Sun Bear Trade in Sarawak, Malaysia." *Traffic Bulletin* 26: 37–40.
- Laurance, W. F., M. Goosem, and S. G. Laurance. 2009. "Impacts of Roads and Linear Clearings on Tropical Forests." *Trends in Ecology & Evolution* 24: 659–669.
- Lee, D. C., V. J. Powell, and J. A. Lindsell. 2019. "Understanding Landscape and Plot-Scale Habitat Utilisation by Malayan Sun Bear (*Helarctos Malayanus*) in Degraded Lowland Forest." *Acta Oecologica* 96: 1–9.
- Lee, S. X. T., Z. Amir, J. H. Moore, K. M. Gaynor, and M. S. Luskin. 2024. "Effects of Human Disturbances on Wildlife Behaviour and Consequences for Predator-Prey Overlap in Southeast Asia." *Nature Communications* 15: 1521.
- Li, F., X. Zheng, X.-L. Jiang, and B. P. L. Chan. 2017. "Rediscovery of the Sun Bear (*Helarctos malayanus*) in Yingjiang County, Yunnan Province, China." *Zoological Research* 38: 206.
- Lieb, Z. E., E. Meijaard, J. F. Brodie, et al. 2025. "Mapping Multiple Wild Pig Species' Population Dynamics in Southeast Asia During the African Swine Fever Outbreak (2018–2024)." *Conservation Letters* 18: e13105.
- Linkie, M., Y. Dinata, A. Nugroho, and I. A. Haidir. 2007. "Estimating Occupancy of a Data Deficient Mammalian Species Living in Tropical Rainforests: Sun Bears in the Kerinci Seblat Region, Sumatra." *Biological Conservation* 137: 20–27.
- Linkie, M., D. J. Martyr, A. Harihar, et al. 2015. "Editor's Choice: Safeguarding Sumatran Tigers: Evaluating Effectiveness of Law Enforcement Patrols and Local Informant Networks." *Journal of Applied Ecology* 52: 851–860.

- Luke, S. H., T. M. Fayle, P. Eggleton, E. C. Turner, and R. G. Davies. 2014. "Functional Structure of Ant and Termite Assemblages in Old Growth Forest, Logged Forest and Oil Palm Plantation in Malaysian Borneo." *Biodiversity and Conservation* 23: 2817–2832.
- Luskin, M. S., L. Arnold, A. Sovie, et al. 2023a. "Mesopredators in Forest Edges." *Wildlife Letters* 1: 107–118.
- Luskin, M. S., J. S. Brashares, K. Ickes, et al. 2017. "Cross-Boundary Subsidy Cascades From Oil Palm Degrade Distant Tropical Forests." *Nature Communications* 8: 2231.
- Luskin, M. S., E. D. Christina, L. C. Kelley, and M. D. Potts. 2014. "Modern Hunting Practices and Wild Meat Trade in the Oil Palm Plantation-Dominated Landscapes of Sumatra, Indonesia." *Human Ecology* 42: 35–45.
- Luskin, M. S., J. S. H. Lee, D. P. Edwards, L. Gibson, and M. D. Potts. 2018. "Study Context Shapes Recommendations of Land-Sparing and Sharing; a Quantitative Review." *Global Food Security* 16: 29–35.
- Luskin, M. S., E. Meijaard, S. Surya, I. Sheherazade, C. Walzer, and M. Linkie. 2021. "African Swine Fever Threatens Southeast Asia's 11 Endemic Wild Pig Species." *Conservation Letters* 14: e12784.
- Luskin, M. S., J. H. Moore, C. P. Mendes, M. B. Nasardin, M. Onuma, and S. J. Davies. 2023b. "The Mass Mortality of Asia's Native Pigs Induced by African Swine Fever." *Wildlife Letters* 1: 8–14.
- Luskin, M. S., and M. D. Potts. 2011. "Microclimate and Habitat Heterogeneity Through the Oil Palm Lifecycle." *Basic and Applied Ecology* 12: 540–551.
- Mahmood, T., T. T. Vu, A. Campos-Arceiz, et al. 2021. "Historical and Current Distribution Ranges and Loss of Mega-Herbivores and Carnivores of Asia." *PeerJ* 9: e10738.
- Meijaard, E. 1999. "Human-Imposed Threats to Sun Bears in Borneo." *Ursus* 11: 185–192.
- Mendes, C. P., W. R. Albert, Z. Amir, et al. 2024a. "Camtrapasia: A Dataset of Tropical Forest Vertebrate Communities From 239 Camera Trapping Studies." *Ecology* 105: e4299.
- Mendes, C. P., X. Liu, Z. Amir, J. H. Moore, and M. S. Luskin. 2024b. "A Multi-Scale Synthesis of Mousedeer Habitat Associations in Southeast Asia Reveals Declining Abundance but Few Extirpations in Fragments and Edges." *Austral Ecology* 49: e13470.
- Mendes, C. P., and M. S. Luskin. 2025. "The Use, Misuse and Opportunities for Structural Equation Modelling (SEM) in Wildlife Ecology." *Journal of Applied Ecology* 62: 3210–3226.
- Miettinen, J., C. Shi, and S. C. Liew. 2016. "2015 Land Cover Map of Southeast Asia at 250 m Spatial Resolution." *Remote Sensing Letters* 7, no. 7: 701–710. <https://doi.org/10.1080/2150704x.2016.1182659>.
- Nazeri, M., L. Kumar, K. Jusoff, and A. R. Bahaman. 2014. "Modeling the Potential Distribution of Sun Bear in Krau Wildlife Reserve, Malaysia." *Ecological informatics* 20: 27–32.
- Negret, P. J., M. S. Luskin, B. Gomez-Valencia, et al. 2023. "Neotropical Understory Birds and Mammals Show Divergent Behaviour Responses to Human Pressure." *Perspectives in ecology and conservation* 21: 180–188.
- Normua, F., S. Higashi, L. Ambu, and M. Mohamed. 2004. "Notes on Oil Palm Plantation Use and Seasonal Spatial Relationships of Sun Bears in Sabah, Malaysia." *Ursus* 15: 227–231.
- Nursamsi, I., Z. Amir, H. Decoeur, J. H. Moore, and M. S. Luskin. 2023. "Sunda Pangolins Show Inconsistent Responses to Disturbances Across Multiple Scales." *Wildlife Letters* 1: 59–70.
- O'Kelly, H. J., J. M. Rowcliffe, S. M. Durant, and E. J. Milner-Gulland. 2018. "Robust Estimation of Snare Prevalence Within a Tropical Forest Context Using N-Mixture Models." *Biological Conservation* 217: 75–82.
- Rayan, D. M., and M. Linkie. 2020. "Managing Threatened Ungulates in Logged-Primary Forest Mosaics in Malaysia." *PLOS ONE* 15: e0243932.
- Ridout, M. S., and M. Linkie. 2009. "Estimating Overlap of Daily Activity Patterns From Camera Trap Data." *Journal of Agricultural, Biological, and Environmental Statistics* 14: 322–337.
- Ripple, W. J., G. Chapron, J. V. LÓPez-Bao, et al. 2016. "Saving the World's Terrestrial Megafauna." *Bioscience* 66: 807–812.
- Ripple, W. J., C. Wolf, T. M. Newsome, et al. 2019. "Are We Eating the World's Megafauna to Extinction?" *Conservation Letters* 12: e12627.
- Rovero, F., and F. Zimmermann. 2016. *Camera Trapping for wildlife research*. Pelagic Publishing.
- Rowcliffe, J. M., R. Kays, B. Kranstauber, C. Carbone, and P. A. Jansen. 2014. "Quantifying Levels of Animal Activity Using Camera Trap Data." *Methods in Ecology and Evolution* 5: 1170–1179.
- Royle, J. A., and J. D. Nichols. 2003. "Estimating Abundance From Repeated Presence-Absence Data or Point Counts." *Ecology* 84: 777–790.
- Sartor, C. C., Z. Kaszta, J. Kamler, et al. 2024. "Identifying Remnant Biodiversity Hotspots in Southern Asia Reveals Disequilibrium in Mammalian Communities." *Biodiversity and Conservation* 33: 3057–3074.
- Scheffers, B. R., B. F. Oliveira, I. Lamb, and D. P. Edwards. 2019. "Global Wildlife Trade Across the Tree of Life." *Science* 366: 71–76.
- Scotson, L. 2019. "Exploring Potential Range Connectivity of Sun Bear (Carnivora: Ursidae: Ursinae)." *Raffles Bulletin of Zoology* 67: 67–76.
- Scotson, L., G. Fredriksson, D. Augeri, C. Cheah, D. Ngoprasert, and W. Wai-Ming. 2017. *Helarctos malayanus*. The IUCN Red List of Threatened Species 2017:e. T9760A45033547.
- Scotson, L., G. Fredriksson, D. Ngoprasert, W.-M. Wong, and J. Fieberg. 2017. "Projecting Range-Wide Sun Bear Population Trends Using Tree Cover and Camera-Trap Bycatch Data." *PLoS One* 12: e0185336.
- Shahanan, M., and I. Debski. 2002. "Vertebrates of Lambir Hills National Park, Sarawak, Malaysia." *Malayan Nature Journal* 56, no. 1: 103–118.
- Shepherd, C., and L. Shepherd. 2010. "The Poaching and Trade of Malayan Sun Bears in Peninsular Malaysia." *Traffic Bulletin* 23: 49–52.
- Sollmann, R., A. Mohamed, H. Samejima, and A. Wilting. 2013. "Risky Business or Simple Solution – Relative Abundance Indices From Camera-Trapping." *Biological Conservation* 159: 405–412.
- Steinmetz, R., D. L. Garshelis, W. Chutipong, and N. Seuaturien. 2011. "The Shared Preference Niche of Sympatric Asiatic Black Bears and Sun Bears in a Tropical Forest Mosaic." *PLoS One* 6: e14509.
- Strang, K., and N. Rusli. 2021. "The Challenges of Conserving Biodiversity: A Spotlight on Southeast Asia." In *Wildlife Biodiversity Conservation: Multidisciplinary and Forensic Approaches*, edited by S. C. Underkoffler and H. R. Adams, 47–66. Springer International Publishing.
- Tee, T. L., F. T. van Manen, and P. Kretzschmar, et al. 2021. "Anthropogenic Edge Effects in Habitat Selection by Sun Bears in a Protected Area." *Wildlife Biology* 1: 00776–00711.
- UNEP-WCMC and IUCN. 2021. *Protected Planet: The World Database on Protected Areas (WDPA)*. UNEP-WCMC and IUCN, Cambridge.
- Wilcove, D. S., X. Giam, D. P. Edwards, B. Fisher, and L. P. Koh. 2013. "Navjot's Nightmare Revisited: Logging, Agriculture, and Biodiversity in Southeast Asia." *Trends in Ecology & Evolution* 28: 531–540.
- Williams, P. J., R. C. Ong, J. F. Brodie, and M. S. Luskin. 2021. "Fungi and Insects Compensate for Lost Vertebrate Seed Predation in An Experimentally Defaunated Tropical Forest." *Nature Communications* 12: 1650.
- Te Wong, S., C. Servheen, L. Ambu, and A. Norhayati. 2005. "Impacts of Fruit Production Cycles on Malayan Sun Bears and Bearded Pigs in Lowland Tropical Forest of Sabah, Malaysian Borneo." *Journal of Tropical Ecology* 21: 627–639.

Te Wong, S., C. W. Servheen, and L. Ambu. 2004. "Home Range, Movement and Activity Patterns, and Bedding Sites of Malayan Sun Bears *Helarctos Malayanus* in the Rainforest of Borneo." *Biological Conservation* 119: 169–181.

Wong, S. T. 2002. "Food Habits of Malayan Sun Bears in Lowland Tropical Forest of Borneo." *Ursus* 13: 127–136.

Wong, W. M., N. Leader-Williams, and M. Linkie. 2013. "Quantifying Changes in Sun Bear Distribution and Their Forest Habitat in Sumatra." *Animal Conservation* 16: 216–223.

Wong, W. M., and M. Linkie. 2013. "Managing Sun Bears in a Changing Tropical Landscape." *Diversity and Distributions* 19: 700–709.

### **Supporting Information**

Additional supporting information can be found online in the Supporting Information section.

Sun Bear Supporting Information 20260105.